

Research Article

Identification of Potential Pathogen Bacteria Causing Tuber Rot in Porang (*Amorphophallus muelleri* Blume)

Afifah Nur Aini, Rodiyati Azrianingsih *, Irfan Mustofa

Department of Biology, University of Brawijaya, Malang 65145, Indonesia

Article history:

Submission April 2019

Revised August 2019

Accepted November 2019

*Corresponding author:

E-mail: rodiyati@ub.ac.id

ABSTRACT

Tuber rot has been reported as one of the limiting factors to porang (*Amorphophallus muelleri* Blume) tuber production. Bacterial infection is one of the main causes of tuber rot during storage. In this study, a total of seven bacterial isolates obtained from porang tubers were tested for their potency to cause rotting on porang tubers. Porang tubers were immersed in bacterial suspension and incubated for 14 days. The physical properties of the tubers such as texture and color, and the chemical properties such as reducing sugar and ethanol content were determined at the end of incubation period. All infected tubers were soft, slimy and darker. Two isolates, T4 and T9A, had the highest potency to cause tuber rot indicated by the significantly higher reducing sugar content (14.38 and 15.32 mg/ml) and higher ethanol content (32.17 and 35.05%) compared to control tubers which were not infected with bacteria. Both isolates were identified as *Bacillus altitudinis* and *Pseudomonas stutzeri*, respectively, based on 16S rDNA sequence analysis.

Keywords: *Amorphophallus muelleri*, pathogen bacteria, tuber rot, porang

Introduction

Porang (*Amorphophallus muelleri* Blume) is a perennial tuberous herb belonging to the Araceae family. *A. muelleri* tuber is rich in glucomannan (up to 70%) [1], a polysaccharide with many industrial uses and health benefits. This property makes porang tuber a valuable commodity. Every year, Indonesia exports around 300 tons of *porang* dried chips to Japan, China and Korea [2].

The tuber production, however, is hindered by the decreasing quality and quantity of tubers due to tuber rot. Freshly harvested *porang* tuber has high water content (75–87%) [3], which provides a favorable environment for the growth of rot-causing microorganisms. The climate in Indonesia which has warm temperature and high humidity is stated to have a role in promoting soft rot disease [4].

Some of the bacteria that have been reported as the cause of tuber rot (also known as soft rot bacteria) in *Amorphophallus* spp. are *Pectobacterium chrysanthemi* [5], *Dickeya dadantii* [6], and

sometimes saprophytic bacteria such as *Bacillus*, *Pseudomonas* and *Clostridium* spp. [7]. Bacterial infection of tubers occurs through the wound caused by harvesting or insect bites.

Tubers infected with soft rot bacteria undergo physical and chemical changes. The rotting tubers become soft and slimy, change color from yellow to dark brown or black, and emit an unpleasant smell [4]. The severity of soft rot disease had a positive correlation with the relative total sugars and reducing sugars in potatoes [4]. Volatile metabolites, such as methanol, ethanol, propanol and butanol, were produced higher in potatoes that are infected with soft rot bacteria than in healthy ones [8].

Exploration of new pathogen species is important because the information will be useful in determining tuber rot control methods and improving the quality and quantity of tuber. Therefore, this study is intended to determine the bacteria isolated from *porang* tuber that have the high-

How to cite:

Aini NA, Azrianingsih R, Mustofa I (2020) Identification of Potential Pathogen Bacteria Causing Tuber Rot in Porang (*Amorphophallus muelleri* Blume). Journal of Tropical Life Science 10 (2): 99 – 104. doi: 10.11594/jtls.10.02.02.

est potency to cause rotting using physical and chemical properties of the tuber as parameters and to identify it based on 16S rDNA sequence.

Material and Methods

There is a total of seven bacterial isolates that were obtained from the collection of Microbiology Laboratory, Faculty of Mathematics and Natural Sciences, University of Brawijaya. The isolates were originally isolated by previous researchers from porang tubers cultivated in Rejosari Village, Bantur, Malang. The isolation was done during storage period of *porang* tubers.

Bacterial culture preparation

Bacterial isolates were sub-cultured using streak and spread method on NA (Nutrient Agar) media and incubated at 30°C for 24 h. The isolates were purified based on colony morphology and confirmed with Gram staining. If different colonies were found, re-purification was carried out until single colony was obtained. The isolates were cultured on slant agars and stored at 4°C for future needs [9].

Pathogenicity test

Pathogenicity test was based on Koch's Postulate principle. Each isolate was cultured in NB (Nutrient Broth) media at 30°C for 24 h. After the cell density of each culture was equalized, bacterial suspension was made by inoculating 2 ml of bacterial culture into 20 ml of sterile NB in a culture bottle closed with cotton plug. *Porang* tuber slices were surface-sterilized by soaking in 5% NaOCl for two minutes and rinsing with sterile water for one minute twice [10]. The tuber slices (triplicate) were immersed in the prepared bacterial suspension of each isolate and incubated at 30°C for 14 days [11]. At the end of incubation period, each tuber slice was taken out of the suspension and put in a Petri dish to observe the color

Sample Preparation for Tuber Chemical Assays
Each tested tuber slice (2 g) was thoroughly pulverized in a mortar and put in a centrifuge tube. The bacterial suspension (10 ml) from which the tuber slice was taken was added into the same tube. The mixture was centrifuged at 5.000 rpm, 10°C for 10 minutes. Supernatant was used for the determination of reducing sugar and ethanol content.

Reducing sugar assay

Reducing sugar content in tuber was determined using spectrophotometric method [12]. The prepared supernatant (1.5 ml) was added with 1.5 ml of distilled water and 3 ml of DNS (dinitrosalicylic acid) reagent in a test tube. The mixture was vortexed and heated in a waterbath at 90°C for 10 minutes. After the mixture was cooled down, 1 ml of 40% Rochelle salt solution was added. The absorbance of the mixture was measured at 540 nm. Reducing sugar content was calculated from a standard curve made by using 2.5, 5, 7.5, 10, 12.5 and 15 mg/ml glucose solution.

Ethanol assay

Ethanol content in tuber was determined using spectrophotometric method [13]. The prepared supernatant (3 ml) was added with 1.5 ml of potassium dichromate ($K_2Cr_2O_7$) reagent in a capped test tube. The absorbance of the mixture was measured at 595 nm. Ethanol content was calculated from a standard curve made by using 2.5, 5, 7.5, 10, 12.5 and 15% ethanol.

Data analysis

The data from tuber chemical assays were analyzed with One Way Analysis of Variance (ANOVA) using IBM SPSS Statistics 20 program. Significant difference is indicated with p-value less than 0.05.

Bacterial identification

The isolate with the highest potency to cause rotting was selected for molecular identification. Bacterial DNA was extracted following the protocol from Zymo Research's Quick-DNA™ Miniprep Kit. The 16S rDNA sequence was amplified by PCR using the conserved primers 27f (5'-AGAGTTTGATCCTGGCTAG-3') and 1492r (5'-CTACGGCTACCTTGTACGA-3'). PCR thermal conditions were as follows: predenaturation at 94°C for 5 min, denaturation at 94°C for 30 s, annealing at 55°C for 30 s, extension at 72°C for 90 s and post extension at 72°C for 5 min [14]. Forward and reverse strands of the amplified DNA fragment were sequenced in an Automatic Sequencer Analyzer ABI 3130 sequencer (1st Base, Malaysia). The forward and reverse sequences were edited using Sequence Scanner v1.0 program and combined using BioEdit v7.2.5 program. The




Figure 1. The condition of tubers after 14 days of incubation. Control tuber (a) and tuber infected with bacterial isolate T4 (b) and T9A (c).

16S rDNA sequence of the isolate was aligned together with reference 16S rDNA sequences obtained from NCBI database. The phylogenetic tree was constructed using MEGA 6.00 program [15].

Results and Discussions

Pathogenicity of bacterial isolates on porang tubers

A total of seven isolates from laboratory collection were tested for the potency to cause rotting on *porang* tubers. Each isolate was shown to have pathogenic properties based on the physical parameters. There were texture and color changes in tubers infected with bacteria and control tuber which was not infected with bacteria.

After 14 days of incubation, all tubers infected with bacteria had soft and slimy texture and darker color, while the control tubers had rigid texture and light-yellow color. The most notable changes were observed in tubers infected with isolate T4 and T9A in which the texture was the softest and the color changed into brownish yellow and light brown (Figure 1).




Figure 2. Reducing sugar in tubers infected with bacterial isolates after 14 days of incubation

The texture change in rotting tuber is related to the tissue maceration caused by the pectinolytic enzymes secreted by pathogenic bacteria. Soft rot bacteria secreted pectinolytic enzymes that break down pectin in the middle lamella, that acts as connector between cells and gives rigidity [16]. The pectinolytic enzymes include pectinases, cellulases, proteases, and xylanases [7]. The color change in rotting tuber due to necrosis. After the cell wall components degrade, the dead tissue will dry up and turn color into light brown or beige [17].

Despite the physical properties of all infected tubers showing symptoms of rotting, the pathogenicity of bacterial isolates should also be determined by their ability to alter the chemical contents in tuber. The reducing sugar and ethanol assay of tuber was done after 14 days of incubation. Based on the reducing sugar assay, there were two isolates showing the highest reducing sugar content out of all tubers tested (Figure 2). The reducing sugar content in tubers infected with isolate T4 and T9A were 14.38 and 15.32 mg/ml, respectively. These two isolates were not statistically different, but the difference between them and the other isolates was significant ($p < 0.05$).

Isolate T4 and T9A also had the highest ethanol content out of all tubers tested in the ethanol assay with ethanol percentage of 32.17 and 35.05%, respectively (Figure 3). There was no significant difference between these two isolates, but they were statistically higher than the other isolates ($p < 0.05$). Based on these results, isolate T4 and T9A were selected as the potential pathogen bacteria with the highest potency to cause rotting on *porang* tuber.




Figure 3. Ethanol in tubers infected with bacterial isolates after 14 days of incubation.




Figure 4. Neighbor-joining phylogenetic tree based on 16S rDNA showing the distance of T4 with reference isolates

The chemical changes in rotting tuber is due to active metabolism of pathogen bacteria. Bacteria secrete amylase to break down the starch in tuber [18], of which a fresh porang tuber contains 15.7% [19]. Starch is broken down to maltose, maltose subsequently is hydrolyzed to glucose.

Glucose is further metabolized by bacteria into pyruvic acid through glycolysis. The pyruvic acid can be either metabolized aerobically or anaerobically. The aerobic pathway ends in TCA cycle, whereas the anaerobic pathway ends in the conversion of acetaldehyde to ethyl alcohol or ethanol [18]. These explanations support the result of this study that the reducing sugar and ethanol content is higher in tubers infected with bacteria than in control tubers (C) (Figure 2 and 3).

Even so, just like other plant materials, tubers




Figure 5. Neighbor-joining phylogenetic tree based on 16S rDNA showing the distance of isolate T9A with reference isolates.

naturally contain sugars such as fructose, sucrose, cellobiose, rhamnose and mannitol [18]. This explains the reducing sugars present in control tubers (C) (Figure 2). It is possible that the ethanol present in control tubers (C) (Figure 3) was due to the low amount of oxygen inside the culture bottle during incubation. There has been a report of endogenous ethanol accumulation in potato tubers under anaerobic condition [20].

Identification of potential pathogen bacteria

Isolate T4 had 99.7% 16S rDNA sequence similarity with *Bacillus altitudinis* strain 41KF2b (Figure 4). Isolate T9A had 99.1% 16S rDNA sequence similarity with *Pseudomonas stutzeri* strain ATCC 17588 (Figure 5). Genus *Bacillus* and *Pseudomonas* are known to be saprophytic bacteria that are commonly present on tubers [7]. *Bacillus* and *Pseudomonas* are some of the dominant groups accompanying *Pectobacterium atrosepticum* during the rotting of potato tubers [21]. Both genera are known as commensals, non-pathogenic microbes, that invade the intracellular space of the host, but are usually dormant. However, once the tubers are infected with pectinolytic bacteria like *Pectobacterium* spp., they began to flourish during the rotting process [21].

So far, *B. altitudinis* has been reported as a soft rot agent in apples and pears [22]. Although the information about pathogenicity of *B. altitudinis* is still limited, there are several reports of other closely-related *Bacillus* species as soft rot agent,

such as *B. pumilus* in potato [23], ginger [24] and melon [25], *B. polymyxa* in garlic [26], and *B. amyloliquefaciens* in potato [27].

P. stutzeri isolated from spoilt orange has the ability to produce pectinase [28]. *P. stutzeri* has a relatively high conversion rate of starch [29]. This explains the high reducing sugar content in tubers infected with isolate T9A. *P. stutzeri* is known to cause brown spot disease in oyster mushroom [30]. Other *Pseudomonas* species that could cause soft rot, such as *P. cichorii* and *P. maculicola* in cruciferous vegetables [18], *P. marginalis* in potato [31] and tuber of *Zantedeschia* spp. which are from the same family as porang [32], *P. aeruginosa* in onion [33] and ginseng [34], and *P. viridiflava* in tomato [35].

Conclusion

Bacterial isolates from *porang* tubers that have the highest potency to cause rotting on *porang* tubers were T4 and T9A, which were identified as *Bacillus altitudinis* and *Pseudomonas stutzeri*, respectively. Both genera have the ability to break down tuber starch into sugars and ferment sugars into ethanol and act as opportunistic pathogen of tuber rot.

Acknowledgment

This research was financially supported by DPPM/DIKTI Research Grant 2018, Faculty of Mathematics and Natural Sciences, Brawijaya University.

References

- Mekkerdchoo O, Borompichaichartkul C, Perrigo A et al. (2016) Tracing the Evolution and Economic Potential of Konjac Glucomannan in *Amorphophallus* species (Araceae) using Molecular Phylogeny and RAPD Markers. *Phytotaxa* 282 (2): 81. doi: 10.11646/phytotaxa.282.2.1.
- Santosa E, Susila AD, Lontoh AP et al. (2016) NPK Levels and Application Methods on Productivity of *Amorphophallus muelleri* Blume in Intercropping System. *Jurnal Jamu Indonesia* 1 (2): 1–8. doi: 10.29244/jji.v1i2.12.
- Dwiyono K, Sunarti TC, Suparno O, Haditjaroko L (2012) Patchouli Alcohol Enrichment From Patchouli Oil Using Molecular Distillation Unit. *Journal of Agroindustrial Technology* 24 (3): 179–188.
- Nourian F, Kushalappa AC, Ramaswamy HS (2002) Physical, Physiological and Chemical Changes in Potato as Influenced By *Erwinia Carotovora* Infection. *Journal of Food Processing and Preservation* 26 (5): 339–359. doi: 10.1111/j.1745-4549.2002.tb00489.x.
- Wu J, Yang C, Jiao Z et al. (2015) Genetic Relationships of Soft Rot Bacteria Isolated from Konjac in China by Amplified Fragment Length Polymorphism (AFLP) and 16S rDNA Gene Sequences. *Agricultural Sciences* 06 (07): 717–723. doi: 10.4236/as.2015.67069.
- He F, Duan J, Luo B et al. (2013) Pathogenic bacteria biodiversity of *Amorphophallus* infected with soft rot disease in Langao, Shaanxi. *Journal of Northwest A & F University (Natural Science Edition)* 41 (7): 91–98.
- Pérombelon MCM (2002) Potato diseases caused by soft rot erwinias: an overview of pathogenesis. *Plant Pathology* 51 (1): 1–12. doi: 10.1046/J.0032-0862.2001.SHORTTITLE.DOC.X.
- Waterer DR, Pritchard MK (1985) Production of volatile metabolites in potatoes infected by *erwinia carotovora* var. *carotovora* and *e. carotovora* var. *atroseptica*. *Canadian Journal of Plant Pathology* 7 (1): 47–51. doi: 10.1080/07060668509501513.
- Kesaulya H, Baharuddin, Zakaria B, Syaiful SA (2015) Isolation and Physiological Characterization of PGPR from Potato Plant Rhizosphere in Medium Land of Buru Island. *Procedia Food Science* 3 190–199. doi: 10.1016/J.PROFOO.2015.01.021.
- Arfani N, Azrianingsih R, Suharjo S, Suharjo S (2018) Isolation and Identification of Antagonistic Bacterium against a Pathogens of Bacterial Tuber Rot of *Amorphophallus muelleri*. *The Journal of Experimental Life Sciences* 8 (3): 165–172. doi: 10.21776/ub.jels.2018.008.03.06.
- Sakaroni R, Suharjo S, Azrianingsih R (2019) Identification of potential pathogen fungi which cause rotten on *Porang (Amorphophallus muelleri* Blume) tubers. In: *AIP Conf. Proc.* AIP Publishing LLC . p 080010.
- Ikanone CEO, Oyekan PO (2014) Effect of Boiling and Frying on the Total Carbohydrate, Vitamin C and Mineral Contents of Irish (*Solanum tuberosum*) and Sweet (*Ipomea batatas*) Potato Tubers. *Nigerian Food Journal* 32 (2): 33–39. doi: 10.1016/S0189-7241(15)30115-6.
- Sayyad SA, Chaudhari SR, Panda BP (2015) Quantitative determination of ethanol in arishta by using UV-visible spectrophotometer. *Pharmaceutical and Biological Evaluations* 2 (5): 204–207.
- Naveed M, Mubeen S, Khan S et al. (2014) Identification and characterization of rhizospheric microbial diversity by 16S ribosomal RNA gene sequencing. *Brazilian Journal of Microbiology* 45 (3): 985–993. doi: 10.1590/S1517-83822014000300031.

15. Mohkam M, Nezafat N, Berenjian A et al. (2016) Identification of *Bacillus* Probiotics Isolated from Soil Rhizosphere Using 16S rRNA, recA, rpoB Gene Sequencing and RAPD-PCR. *Probiotics and Antimicrobial Proteins* 8 (1): 8–18. doi: 10.1007/s12602-016-9208-z.
16. Wakil SM, Oyinlola KA (2011) Diversity of pectinolytic bacteria causing soft rot disease of vegetables in Ibadan, Nigeria. *Journal of Applied Biosciences* 38 2540–2550.
17. Benada M, Boumaaza B, Boudalia S et al. (2018) Variability of aggressiveness and virulence of *Erwinia carotovora* subsp. *carotovorum* causing the soft rot on potato tubers in the western of Algeria. *Int J Plant Biol.* doi: 10.4081/pb.2018.7568
18. Tournas VH (2005) Spoilage of Vegetable Crops by Bacteria and Fungi and Related Health Hazards. *Critical Reviews in Microbiology* 31 (1): 33–44. doi: 10.1080/10408410590886024.
19. Rokhmah DN, Supriadi H (2015) Prospek Pengembangan Iles-Iles (*Amorphophallus muelleri* Blume) sebagai Upaya Diversifikasi Pangan di Indonesia. *SIRI-NOV* 3 (1): 1–10.
20. Claassens MMJ, Verhees J, van der Plas LHW et al. (2005) Ethanol breaks dormancy of the potato tuber apical bud. *Journal of Experimental Botany* 56 (419): 2515–2525. doi: 10.1093/jxb/eri245.
21. Kõiv V, Roosaare M, Vedler E et al. (2015) Microbial population dynamics in response to *Pectobacterium atrosepticum* infection in potato tubers. *Scientific Reports* 5 (1): 11606. doi: 10.1038/srep11606.
22. Elbanna K, Elnaggar S, Bakeer A (2014) Characterization of *Bacillus altitudinis* as a New Causative Agent of Bacterial Soft Rot. *Journal of Phytopathology* 162 (11–12): 712–722. doi: 10.1111/jph.12250.
23. Bathily H, Babana AH, Samaké F (2010) *Bacillus pumilus*, a new pathogen on potato tubers in storage in Mali. *African Journal of Microbiology Research* 4 (20): 2067–2071.
24. Peng Q, Yuan Y, Gao M (2013) *Bacillus pumilus*, a Novel Ginger Rhizome Rot Pathogen in China. *Plant Disease* 97 (10): 1308–1315. doi: 10.1094/PDIS-12-12-1178-RE.
25. Song JH, Wu ZR, Zhang LX et al. (2018) First Report of *Bacillus pumilus* Causing Fruit Rot on Muskmelon (*Cucumis melo*) in China. *Plant Disease* 102 (2): 439–439. doi: 10.1094/PDIS-08-17-1169-PDN.
26. Galal AA, Abdel-Gawad TI, El-Bana AA (2002) Post-harvest decay of garlic cloves caused by *Bacillus polymyxa* and *Fusarium moniliforme*. *Egypt Journal of Microbiology* 36: 71–88.
27. Wang L, Li XB, Suo HC et al. (2017) Soft rot of potatoes caused by *Bacillus amyloliquefaciens* in Guangdong province, China. *Canadian Journal of Plant Pathology* 39 (4): 533–539. doi: 10.1080/07060661.2017.1381994.
28. Muslim DSN, Hasan AM, Mahdi NZ (2016) Antibiofilm and antiadhesive properties of pectinase purified from *Pseudomonas stutzeri* isolated from spoilt orange. *Advances in Environmental Biology* 10 (11): 91–99.
29. Schmidt J, John M (1979) Starch metabolism in *Pseudomonas stutzeri*. I. Studies on maltotetraose-forming amylase. *Biochimica et Biophysica Acta (BBA) - Enzymology* 566 (1): 88–99. doi: 10.1016/0005-2744(79)90252-3.
30. Mallesha BC, Shetty KS (1988) A New Brown Spot Disease of Oyster Mushroom Caused by *Pseudomonas stutzeri*. *Current Science* 57 1190–1192. doi: 10.2307/24090888.
31. Li J, Chai Z, Yang H et al. (2007) First report of *Pseudomonas marginalis* pv. *marginalis* as a cause of soft rot of potato in China. *Australasian Plant Disease Notes* 2 (1): 71. doi: 10.1071/DN07029.
32. Krejzar V, Mertelík J, Pánková I et al. (2008) *Pseudomonas marginalis* associated with soft rot of *Zantedeschia* spp. *Plant Protection Science* 44: 85–90. doi: 10.17221/16/2008-PPS.
33. Abd-Alla MH, Bashandy SR, Ratering S, Schnell S (2011) First report of soft rot of onion bulbs in storage caused by *Pseudomonas aeruginosa* in Egypt. *Journal of Plant Interactions* 6 (4): 229–238. doi: 10.1080/17429145.2010.535618.
34. Gao J, Wang Y, Wang CW, Lu BH (2014) First report of bacterial root rot of ginseng caused by *Pseudomonas aeruginosa* in China. *Plant Disease* 98 (11): 1577. doi: 10.1094/PDIS-03-14-0276-PDN.
35. Saygili H, Sahin F, Sahin F et al. (2005) New symptoms of tomato soft rot diseases in Turkey. *Acta Horticulturae* 695: 291–294. doi: 10.17660/ActaHortic.2005.695.32.