

PCR-RFLP and Sequencing of *trnS/trnfM* Fragment of *Enhalus acoroides* from Sanur Coastal Waters, Bali, Indonesia: A Preliminary Study

Made Pharmawati*, Eka Fibayani Imaniar

Biology Department, Faculty of Mathematics and Natural Sciences, Udayana University, Badung, Indonesia

ABSTRACT

This study aimed to evaluate the resolution of PCR-RFLP and sequencing of *trnS/trnfM* in detecting genetic diversity of seagrass *Enhalus acoroides* at Sanur Beach, Sindhu Beach and Semawang Beach. This research used six samples *E. acoroides* from each location. DNA extraction method followed Doyle and Doyle. PCR amplifications were done using primers P672/P673, P674/P675 and ITS4/ITS5 followed by digestion using restriction enzymes *EcoRI*, *PstI*, *HindIII*, *BamHI*, *RsaI*, *MvaI* and *HinfI*. Sequencing analysis used PCR products of primers *trnS/trnfM*. Testing of DNA sequences of *E. acoroides* were conducted using BLAST (Basic Local Alignment Search Tool). DNA sequences were further analyzed using MEGA 5.2 software (Molecular Evolutionary Genetic Analysis) to evaluate variations of DNA. The sequence alignments were done using ClustalW software to determine the homology between the DNA sequences. The results showed that 18 samples of *E. acoroides* from Sanur, Sindhu and Semawang Beach have no polymorphism based on restriction enzyme analyses. Furthermore, sequencing of *trnS/trnfM* region of 18 samples *E. acoroides* showed that the sequences were identical.

Keywords: *Enhalus acoroides*, PCR-RFLP, Sequencing *trnS/trnfM*, Sanur

INTRODUCTION

Seagrass meadow is one of coastal ecosystem with seagrass as dominant vegetation. Seagrass meadow may consist of single seagrass species or mix of two to 12 seagrass species on several types of substrates [1].

Seagrass ecosystem has high productivity through high CO₂ capture and nutrient trap [2]. Seagrass meadow has important functions both ecologically and economically. From ecological point of view, seagrass meadow acts as nursery ground and spawning area for marine biotas such as fish, mollusc, sea urchin, sea star and other marine animals [3]. Besides that, seagrass itself performs as food resources for dugong (*Dugong dugon*) and green turtle (*Chelonia mydas*) [4]. Economically, beach with seagrass bed can be as recreational and educational areas.

Seagrass grows in shallow coastal waters at intertidal and sublittoral zones, in between high and low tide, and around rocky island in Indonesia [5]. There

are 60 seagrass species identified in the world, and 13 of them are found in Indonesia [6].

In 2010, it was reported that there were eight species of seagrass in coastal waters of Denpasar, Bali [7]. A study in 2013 found that there were nine seagrass species in Sanur Beach and Sindhu Beach, Sanur, Denpasar, Bali [8]. One of seagrass species is *Enhalus acoroides* which has the largest and longest leaf blade and can grow up to 100 cm. *Enhalus acoroides* is seagrass species that lives in all types of waters and grows well in shallow open waters [9].

At south east of Bali beaches, especially in Sanur areas, seagrass bed spreads from Sanur Beach to Mertasari Beach [10]. Semawang Beach is located adjacent to Mertasari Beach. Sindhu Beach is located 500 m from Sanur Beach while the distance of Sindhu Beach to Semawang Beach is approximately 4 km. Information on genetic diversity of *E. acoroides* at Sanur waters is useful for monitoring and determination of con-

*Corresponding author:

Made Pharmawati

Biology Department, Faculty of Mathematics and Natural Sciences, Udayana University

Jalan Kampus Bukit Jimbaran, Jimbaran, Kuta Selatan, Kabupaten Badung, Bali, Indonesia 82121

E-mail: pharmawati@hotmail.com

How to cite:

Pharmawati M, Imaniar EF (2016) PCR-RFLP and sequencing of *trnS/trnfM* fragment of *Enhalus acoroides* from Sanur coastal waters, Bali, Indonesia: A preliminary study. J. Trop. Life. Science 6 (2): 106 – 122.

servation strategy.

Genetic diversity can be detected using molecular markers. One of molecular marker available is PCR-RFLP, which can detect diversity in intraspecific levels [11]. This marker is co-dominant marker and can detect high polymorphism [11]. PCR-RFLP technique has been used to discriminate Fagaceae species [12] as well as to detect variation in chloroplast DNA of wild brassica [13]. It also used to determine mode of inheritance of organelle DNA and phylogenetic relationships in plant [14]. One of chloroplast DNA fragment with high evolution rate is *trnS/trnF* [15]. Beside chloroplast genome, nuclear genome also has high evolution level [16].

More sensitive method to detect DNA variation is DNA sequencing. DNA sequencing can be used to identify and determine the function of DNA fragment by comparing DNA sequence of samples to known DNA [17]. DNA sequencing of chloroplast regions has been employed to detect genetic diversity of *Shorea cutisii* [18], *Pistacia vera* L. [19], and to study phylogeny of *Chrysanthemum* [20].

In this study, PCR-RFLP of nuclear and chloroplast DNA were employed to analyse genetic diversity of *E. acoroides* from Sanur Beach, Sindhu Beach and Semawang Beach. Also, sequencing of *trnS/trnF* fragment of chloroplast DNA was done to support the analyses.

MATERIALS AND METHODS

Sample collection

Enhalus acoroides samples were collected from three locations of Sanur waters (Figure 1) including Sanur Beach (S 8°40'16.52", E 115°15'91.33"), Sindhu Beach (S 8°68'51", E 115°26'53.8") and Semawang Beach (S 8°42'52.98", E 115°15'85.69"). Transect line of 200 m long was placed perpendicular to seashore. Two leaf samples were collected at seashore, two samples at the middle of transect line and two samples at the end of transect line. The distance between two samples was at least 5 m. At each sampling site, leaf blade was collected from one individual.

DNA extraction

DNA extraction was conducted using CTAB extraction buffer [21] with modification by increasing EDTA concentration to 50 mM. Leaf of *E. acoroides* (0.1 g) was ground using mortar and pestle and 1 ml extraction buffer (2% CTAB, 100 mM Tris/HCL pH 8, 1.4 M NaCl, 50 mM EDTA pH 8, 0.2% β -mercaptoethanol) was added. The extract was

transferred to microtube and incubated at 65°C for 30 minutes, followed by centrifugation at 14.000 rpm for 10 minutes. Supernatant was transferred to a new tube and added by the same volume of chloroform : isoamylalcohol (24 : 1). After vortexing for 2 minutes, the tubes were centrifuged at 14.000 rpm for 10 minutes. Aquous layer was transferred to a new tube followed by the addition of the same volume of cold isopropanol.

After overnight incubation at -20°C, samples were centrifuged at 14.000 rpm for 5 minutes. DNA pellet was washed with 70% ethanol and sentrifuged for 3 minutes. DNA pellet was air dried and diluted with 100 μ L sterile water. DNA solution was kept in -20°C.

DNA was visualised using 1% agarose gel electrophoresis stained with ethidium bromide [22]. To estimate DNA concentration, lambda DNA of known concentrations were included in the gel. Electrophoresis was run at 100 V for 30 minutes and visualised using UV transilluminator.

DNA amplification

DNA amplifications were done for chloroplast DNA and nuclear DNA fragments. Table 1 shows the sequences of primer pairs used. Total volume of PCR reaction was 20 μ L containing 75 ng DNA template, 1 \times PCR buffer, 2 μ M MgCl₂, 200 μ M dNTP, 1 unit Taq DNA Polymerase (MDBio) and 2 μ M primer. PCR cycles for amplification are as follow: 1) initial denaturation at 95°C for 5 minutes, 2) denaturation at 95°C for 1 minute, 3) annealing at 56°C (depending on primer pair used) for 2 minutes, 4) extension at 72°C for 1 minute 30 seconds, 5) final extension at 72°C for 10 minutes. Cycles 2 to 4 were repeated for 40 cycles.



Figure 1. Sample collection sites

Table 1. Primers used in this study, their sequences and annealing temperatures

Primer	Location	Amplified region	Sequence	Annealing temperature	Reference
P672	chloroplast	<i>trnQ/rps</i>	5'-GCGTGGCCAAGYGGTAAGGC-3'	56°C	Lucas <i>et al.</i>
P673			5' GTTGCTTTYTACCACATCGTTT-3'		(2012) [23]
ITS4	nucleus	ITS4/ITS5	5'-CCCGCCTGACCTGGGGTCGC-3'	54°C	White <i>et al.</i>
ITS5			5'-TAGAGGAAGGAGAAGTCGTAACAA-3'		(1990) [24]
P674	nucleus	ITS5a/ITS4	5'-CCTTATCATTTAGAGGAAGGAG-3'	62°C	Lucas <i>et al.</i>
P675			5'-TCCTCCGCTTATTGATATGC-3'		(2012) [23]
<i>trnS</i>	chloroplast	<i>trnS/trnfM</i>	5'-GAGAGAGAGGGATTCTGAACC-3'	62°C	Demesure <i>et al.</i>
<i>trnfM</i>			5'-CATAACCTTGAGGTCACGGG-3'		(1995) [25]

Visualisation of PCR products was performed using 1% agarose gel electrophoresis at 100 V for 45 minutes and stained with ethidium bromide. Observation was done using UV transilluminator. As size marker, low DNA mass ladder (Invitrogen) was used.

Restriction enzymes digestion

PCR product of primer pair P672/P673, ITS4/ITS5 and P674/P675 were cut with various restriction enzymes including *EcoRI*, *PstI*, *HindIII*, *BamHI*, *Rsa*, *Mva* and *HinfI*. As many as 4 µL PCR product was included in the reaction together with 4.7 µL H₂O, 1× restriction buffer, and 1 unit restriction enzyme. The reaction was done at 37° C for 2 hours. The results were visualized using 1.5% agarose gel electrophoresis. Low DNA mass ladder (Invitrogen) was included in the gel as size marker.

DNA sequencing

PCR products using primer pair *trnS-trnfM* of 18 *E. acoroides* samples were sent to sequencing facility at Berkeley Sequencing Facility, USA. The sequences were analysed using MEGA 5.2 (Molecular Evolutionary Genetic Analysis) for homolog determination between sequences.

RESULTS AND DISCUSSION

The 18 *E. acoroides* samples were successfully amplified using primer pair P672/P673, ITS4/ITS5, P674/P67, and *trnS/trnfM*. No length polymorphism was detected in all PCR products. Therefore, restriction enzyme digestions were performed. PCR products of primer P672/P673, ITS4/ITS5 and P674/P675 were digested using various restriction enzymes. Restriction enzymes used were *Rsa*, *HindIII*, *Mva*, *BamHI*, *PstI*, *HinfI* and *EcoRI*. The sizes of digested DNA fragments were shown in Table 2. Representatives of digested DNA patterns were presented in Figure 2.

Table 2. Summary of digestion using restriction enzyme

Primer	Size of PCR product (bp)	Restriction enzyme	Size of DNA after digestion (bp)
P672 /	1000	<i>PstI</i>	1000
P673	1000	<i>HindIII</i>	1000
	1000	<i>Mva</i>	1000
	1000	<i>BamHI</i>	1000
	1000	<i>Rsa</i>	600+400=1000
	1000	<i>HinfI</i>	400+380+220=1000
	1000	<i>EcoRI</i>	400+380+220=1000
ITS4 /	550	<i>Mva</i>	550
ITS5	550	<i>Rsa</i>	550
	550	<i>Hind III</i>	550
	550	<i>HinfI</i>	330+220=550
P674 /	800	<i>Rsa</i>	800
P675	800	<i>HindIII</i>	800
	800	<i>Mva</i>	800




Figure 2. Patterns of PCR products digested using restriction enzymes. PCR product using primer P672/P673 digested with *HinfI* (A) and *Rsa* (B). PCR product using primer ITS4/ITS5 digested with *HindIII* (C). PCR product using primer P674/P675 digested using *Mva* (D)

CTCGGCCATCTCTCCGAGAGACAATCTCCATTTTATTCCTATCCATAGTACATAGCCATAGGAATCCATATACTA
 ACGATTTGTAGATGCAGCAACCCACCCTAGAGGGGAATCTAAATTTTCGATGTATCCATCTGTATACGGATTACAG
 TATGTATATCCGTACATTTAGGAAGTCCATACGAAATTCCTGCTAAACGCATTTTCATGTCTAAACCGAAATCCAAG
 AAAGAGTGAATGGTAATAAGTTCTAATAAGTTCTTAAGAAATCCATGGGGGGATTCCTGATGAAATCCCTCCACGA
 TGCATTTTCATCCCAATTTGGACTCAGCGAGGGATCCAATGGTATAGTTCATTTGTTAGTAGTTTGGAGGATTACA
 AGCATGATTCTTGCTTTCCAATTGGCTGTTTTTGCATTAATTGCAACCTCATCCATCCTATTGATTAGTGTACC
 GTTGTATTTGCTTCTTCGGATGGTTGGTCAAGTAACAAAAAGATTGTCTTTTCTGGTACATCATATGGATTGGAT
 TAGTCTTTCTGGTAGCTATCCTGAATTCTCTTATCTCTTGAATCTCTTTGGTTTGGTACCGGATCCAAATCTATA
 TACCTTTCTTCTCAAAAATGCGGATTCTGAGACAATGGAAATGAAATCGGACTGAATTTCTATTAGGAGAAT
 CATCAATCCTAATG

Figure 3. The sequence of *trnS/trnfM* fragment of *E. acoroides*

Restriction enzyme *Pst*I, *Hind*III, *Mva*, *Bam*HI did not cut PCR product of P672/P673. PCR products of primer P674/P675 and ITS4/ITS5 were not digested by *Rsa*, *Hind*III and *Mva*. Table 2 shows that using 14 primer-enzyme combinations there was no polymorphism at 18 *E. acoroides* samples from Sanur coastal region. This result indicates that the population of *E. acoroides* in Sanur Coastal region is quite homogeneous. Based on size estimation using agarose electrophoresis, the length of PCR products using primer *trnS/trnfM* was 1500 bp, while from sequencing analysis only 689 bp of DNA was clearly sequenced (Figure 3). Multiple sequence alignment of 18 individual of *E. acoroides* showed identical base sequences.

Identification of the sequence of *trnS/trnfM* was done using Basic Local Alignment Search Tool (BLAST). The sequence had 97% level of similarity to *Phragmipedium schlimii* and 96% level of similarity to *Ravenala madagascariensis* and *Thaumatococcus daniellii*. This may be because there is no DNA sequence of *trnS/trnfM* fragment of *E. acoroides* in the National Center for Biotechnology Information (NCBI) database.

Both PCR-RFLP and sequencing technique failed to detect variation in *E. acoroides* from Sanur Beach, Sindhu Beach and Semawang Beach, Sanur, Bali. Population of *E. acoroides* at Sanur Beach, Sindhu Beach and Semawang Beach are open population and not isolated by any barrier. Based on the geographic condition, it is predicted that there is gene flow between populations. The presence of gene flow leads to genetic connectivity [26].

A study on development of microsatellite markers for *E. acoroides* found that only 4 pairs of primer out of 36 primer pairs detected polymorphism in *E. acoroides* from Li'Angang and Xincungang, Hainan Province, China [27]. Low genetic diversity was also found in seagrass species *Posidonia oceanica*. In this species, very limited sexual recombination was detected

due to clonal propagation [28].

Rhizome extension is a major factor in development of seagrass meadow. In general, the ability of seagrass to form wide seagrass bed is because of its vegetative growth through the growth and dominance of rhizome [29]. It has been stated that propagation by sexual reproduction and establishment of seedling have little contribution to shoot recruitment in development of seagrass bed [30].

Genetic monomorphism in aquatic plants was detected using allozymes and random amplified polymorphic DNA (RAPD) [31]. This indicated the dominance of clonal propagation than sexual propagation [31, 32, 33] as well as the low resolution of the methods used [31]. It was suggested that *E. acoroides* grown at Sanur Beach, Sindhu Beach and Semawang Beach, Bali is probably grown through vegetative and well developed clonal growth. However, the use of PCR-RFLP and sequencing of limited genome regions is not sufficient to draw robust conclusion. By using 18 samples, this preliminary study indicated that PCR-RFLP of chloroplast and nuclear fragments, as well as sequencing of *trnS/trnfM* showed low resolution.

CONCLUSION

PCR-RFLP of chloroplast and nuclear DNA fragments failed to detect polymorphism of *E. acoroides* from Sanur coastal water, Bali, Indonesia. The sequences of *trnS-trnfM* fragment from 18 individual were identical in all samples.

ACKNOWLEDGMENT

The authors thank USAID for partially funding this research with NAS Sub-Grant No. PGA-2000003438.

REFERENCES

1. Kiswara W, Winardi (1999) Sebaran lamun di Teluk Kuta dan Teluk Gerupuk, Lombok. Jakarta, Pusat Penelitian Oseanografi-LIPI.

2. Duarte CM, Sintes T, Marba N (2013) Assessing the CO₂ capture potential of seagrass restoration projects. *J App Ecol.* doi:10.1111/1365-2664.12155.
3. Barbier EB, Hacker SD, Kennedy C et al (2011) The value of estuarine and coastal ecosystem services. *Ecol Monograph* 81(2): 169-193.
4. Andre J, Gyuris E, Lawder IR (2005) Comparison of the diets of sympatric dugongs and green turtles on the Orman Reefs, Torres Strait, Australia. *Wildlife Res.* 32:53-62.
5. Nienhuis PH, Coosen J, Kiswara W (1989) Community structure and biomass distribution of seagrasses and macrofauna in the Flores Sea, Indonesia. *Neth J Sea Res.* 23(2): 197-214.
6. Kuriandewa TE (2009) Tinjauan tentang Lamun di Indonesia. Lokakarya Nasional I Pengelolaan Ekosistem Lamun. Jakarta, Sheraton Media.
7. Yusup DS, Asy'ari H (2010) Komunitas tumbuhan lamun di kawasan perairan sekitar Denpasar. In Proceedings of the Seminar Nasional Biologi: Biodiversitas dan Bioteknologi Sumberdaya Akuatik. Purwokerto. Edited by Prabowo, Ardli ER, Sastranegara MH, Lestari W, Wijayanti G, Nuryanto A.
8. Nurkamila US (2014) Filogenetik lamun di Pantai Sanur dan Shindu berdasarkan penanda kloroplas DNA. Bachelor Thesis. Udayana University, Biology Department.
9. Kiswara W (2004) Kondisi padang lamun (seagrass) di Perairan Teluk Banten 1998-2001. Lembaga Penelitian Oseanografi. Jakarta, Indonesian Institute of Science.
10. Arthana IW (2004) Spesies dan kerapatan padang lamun di Pantai Sanur Bali. *Bumi Lestari J Env.* 5(2): 1-11.
11. Outram MA, Jones EE, Jaspers MV et al (2014) Development of a PCR-RFLP method to distinguish species within the *Ilyonectria macrodidyma* complex. *New Zealand Pl Protec.* 67: 151-156.
12. Coutinho JP, Carvalho A, José Lima-Brito (2014) Taxonomic and ecological discrimination of Fagaceae species based on ITS PCR-RFLP. *AoB PLANTS plu079* doi:10.1093/aobpla/plu079.
13. Sarin B, Martin JP, Kaula BC, et al (2015) Chloroplast DNA variations in wild Brassicas and their implication in breeding and population genetics studies. *Scientifica* 2015. doi: 10.1155/2015/952395.
14. Pharmawati M, Yan G, Sedgley R et al (2004) Chloroplast DNA inheritance and variation in *Leucadendron* species (Proteaceae) as revealed by PCR-RFLP. *Theor Appl Genet.* 109: 1694-1701.
15. Kress WJ, Wurdack KJ, Zimmer E et al (2005) Use of DNA barcodes to identify flowering plants. *Proc Nat Acad of Sci USA.* 102: 8369-8374.
16. Peay KG, Kennedy PG, Bruns TD (2008) Fungal community ecology: a hybrid beast with a molecular master. *Bio. Sci.* 58: 799-810.
17. Glick BR, Pasternak JJ, Patten CL (2010) *Molecular biotechnology: Principles and applications of recombinant DNA* 4th ed. Washington DC, ASM Press.
18. Kamiya K, Nanami S, Kenzo T et al (2012) Demographic history of *Shorea curtisii* (Dipterocarpaceae) inferred from chloroplast DNA sequence variations. *Biotropica* 44:577-585.
19. Sarra C, Soumaya RC, Zined M et al (2015) Chloroplast DNA analysis of Tunisian pistachio (*Pistacia vera* L.): Sequence variations of the intron trnL (UAA). *Sci Hort.* 191: 57-64.
20. Liu PL, Wan Q, Guo YP et al (2012) Phylogeny of the genus *Chrysanthemum* L.: Evidence from single-copy nuclear gene and chloroplast DNA sequences. *PLoS ONE* 7(11): e-48970. Doi:10.1371/journal.pone.0048970.
21. Doyle JJ, Doyle JL (1990) Isolation of plant DNA from fresh tissue. *Focus* 12:13-15.
22. Sambrook J, Fritsch EF, Maniatis T (1989) *Molecular Cloning: A Laboratory Manual* 2nd Ed. Cold Spring Harbor Laboratory Press, United State America.
23. Lucas C, Tangaradjou T, Papenbrock J (2012) Development of a DNA barcoding system for seagrasses: successful but not simple. *PloS ONE* 7(1): e29987.
24. White TJ, Bruns T, Lee S et al (1990) Amplification and direct sequencing of fungal ribosomal RNA genes for phylogenetics. *New York, Academic Press.*
25. Demesure B, Sodzi N, Petit RJ (1995) A Set of universal primers for amplification of polymorphic non coding regions of mitochondrial and chloroplast DNA in plants. *Mol Ecol.* 4: 129-131.
26. Lowe A, Harris S, Ashton P (2006) *Ecological Genetics (Design, Analysis and Application)*. Singapore, Blackwell publishing.
27. Gao H, Jiang K, Geng Y et al (2012) Development of microsatellite primers of the largest seagrass, *Enhalus acoroides* (Hydrocharitaceae). *Amer J Bot.* e99-e101.
28. Diaz-Almela E, Arnaud-Haond S, Vliet MS et al (2007) Feed-backs between genetic structure and perturbation-driven decline in seagrass (*Posidonia oceanica*) meadows. *Cons Genet.* 8:1377-1391.
29. Waycott M (2000) Matting system and population genetics of marine angiosperm. Wilson, K.L. and Morrison, DA. (eds). *Monocots: Systematics and Evolution*. Melbourne, CSIRO.
30. Lacap CDA, Vermaat JE, Rolton RN et al (2002) Propagule dispersal of the SE Asian seagrasses *Enhalus acoroides* and *Thalassia hemprichii*. *Mar Ecol Prog Ser* 235:75-80.
31. Capiomont A, Sandmeier M, Caye G et al (1996) Enzyme polymorphism in *Posidonia oceanica*, a seagrass endemic

- to the Mediterranean. *Aqua Bot.* 54: 265–277.
32. Procaccini G, Alberte RS, Mazzella L (1996) Genetic structure of the seagrass *Posidonia oceanica* in the western Mediterranean: ecological implications. *Mar Ecol Prog Ser.* 140: 153–160.
 33. Arnaud-Haond S, Alberto F, Teixeira S et al (2005) Assessing genetic diversity in clonal organism: Low diversity or low resolution? Combining power and cost efficiency in selecting markers. *J Hered.* 96: 434-440.